

Differences in life-history and ecological traits between co-occurring *Panulirus* spiny lobsters (Decapoda, Palinuridae)

Patricia Briones-Fourzán¹

¹ *Universidad Nacional Autónoma de México, Instituto de Ciencias del Mar y Limnología, Unidad Académica de Sistemas Arrecifales. Prol. Av. Niños Héroes s/n, Puerto Morelos, Quintana Roo, México*

Corresponding author: Patricia Briones-Fourzán (briones@cmarl.unam.mx)

Academic editor: I.S. Wehrtmann | Received 23 November 2013 | Accepted 29 June 2014 | Published 25 November 2014

<http://zoobank.org/FAC039D4-9EAF-4A9F-9161-C943A82B11F5>

Citation: Briones-Fourzán P (2014) Differences in life-history and ecological traits between co-occurring *Panulirus* spiny lobsters (Decapoda, Palinuridae). In: Wehrtmann IS, Bauer RT (Eds) Proceedings of the Summer Meeting of the Crustacean Society and the Latin American Association of Carcinology, Costa Rica, July 2013. ZooKeys 457: 289–311. doi: 10.3897/zookeys.457.6669

Abstract

Coexistence of closely related species may be promoted by niche differentiation or result from interspecific trade-offs in life history and ecological traits that influence relative fitness differences and contribute to competitive inequalities. Although insufficient to prove coexistence, trait comparisons provide a first step to identify functional differences between co-occurring congeneric species in relation to mechanisms of coexistence. Here, a comparative review on life history and ecological traits is presented for two pairs of co-occurring species of spiny lobsters in the genus *Panulirus*: *P. gracilis* and *P. inflatus* from the Eastern Central Pacific region, and *P. argus* and *P. guttatus* from the Caribbean region. *Panulirus gracilis* and *P. inflatus* have similar larval, postlarval, and adult sizes and a similar diet, but differ in degree of habitat specialization, fecundity, and growth rate. However, little is known on behavioral traits of these two species that may influence their competitive abilities and susceptibility to predators. The more abundant information on *P. argus* and *P. guttatus* shows that these two species differ more broadly in degree of habitat specialization, larval, postlarval and adult sizes, diet, fecundity, growth rate, degree of sociality, defense mechanisms, susceptibility to predators, and chemical ecology, suggesting a greater degree of niche differentiation between *P. argus* and *P. guttatus* than between *P. gracilis* and *P. inflatus*. Whether the substantial niche differentiation and apparent interspecific trade-offs between *P. argus* and *P. guttatus* relative to *P. gracilis* and *P. inflatus* reflect an earlier divergence of the former pair of species in the evolution of the genus constitutes an intriguing hypothesis. However, whether or not post-divergence evolution of each species pair occurred in sympatry remains uncertain.

Keywords

Crustacea, Decapoda, Achelata, Palinuridae, coexistence, resource use, predation, Eastern Central Pacific, Caribbean region

Introduction

Spiny lobsters (Decapoda: Achelata: Palinuridae) are large (body length up to 60 cm), long-lived (> 10 years) crustaceans that occur in a wide range of habitats and depths, and constitute some of the most important fishing resources in all parts of the world (Phillips 2006). Spiny lobsters exhibit complex behaviors and are an important component of community structure and function because they consume a vast array of small benthic organisms and are prey to numerous species of higher predators (Lipcius and Eggleston 2000, Phillips et al. 2013). In addition, spiny lobsters are sturdy and easy to keep in controlled laboratory conditions, making them useful subjects for many types of biological, physiological, and behavioral studies (Cobb 2006).

The family Palinuridae comprises 54 extant species/subspecies arranged in 12 genera (Chan 2010), all of which have a specialized, flat-bodied larva called phyllosoma with multiple stages and a long (4–22 months) planktonic life, as well as a swimming, non-feeding postlarva called puerulus. The most diverse genus is *Panulirus*, with 24 species/subspecies, followed by *Jasus* and *Palinurus*, with six species each. These three genera contain the great majority of commercially important species. However, *Jasus* species are distributed exclusively in cold waters of the southern hemisphere (Jeffs et al. 2013), whereas *Palinurus* species are restricted to south east Africa and the north-eastern Atlantic, and generally occur at depths greater than 100 m (Groeneveld et al. 2013). In contrast, *Panulirus* species occur in shallow tropical and subtropical waters (< 100 m in depth) of both hemispheres, where the diversity of habitats may have promoted a greater radiation of this genus (George and Main 1967, George 2006). Therefore, the occurrence of two or more *Panulirus* species living in sympatry is common in different regions throughout the world (Briones-Fourzán and Lozano-Álvarez 2013).

The co-occurrence of congeneric species at local scales is common in many marine systems (e.g. Azovsky 1996, Sfenthourakis et al. 2005, González et al. 2011), but co-occurrence does not necessarily imply coexistence. The key criterion for coexistence is the “invasibility” criterion, which requires that each species must be able to increase from low density (i.e. persist) when the others are at their typical abundances (Chesson 2000). Conditions that are necessary but not sufficient for invasibility include negative density dependence and trade-offs in performance that influence population regulation (Siepielski and McPeck 2010). Trade-offs imply that advantages that one species may have over others are offset by compensating disadvantages (Chesson 2000, Kneitel and Chase 2004). For example, coexistence of species may be promoted by trade-offs between competitive ability and dispersal ability, between abilities to compete for alternative resources, between competitive ability and disturbance tolerance, and between

competitive ability and susceptibility to predation or disease (Bohannan et al. 2002, Amarasekare 2003, 2008, Kneitel and Perrault 2006). These trade-offs may involve niche differentiation between species (McPeck 1996) or may result from interspecific trade-offs in life history and ecological traits that influence relative fitness differences and contribute to competitive inequalities (e.g. body size, fecundity, longevity, dispersal) (Tilman 1994, Chesson 2000, Amarasekare 2003, Amarasekare et al. 2004).

Because of the wealth of data needed, it is difficult to prove whether co-occurring species truly coexist, particularly for long-lived species wherein the relevant data should have to span multiple generations of each species (Siepielski and McPeck 2010, HilleRisLambers et al. 2012, Narwani et al. 2013). On the other hand, many studies related to coexistence have addressed species belonging to different genera, underscoring the need for more studies focusing on trying to understand the degree to which congeneric species that are within the same trophic level coexist, in particular at local scales (Siepielski and McPeck 2010). As species within genera are often predicted to be more similar to each other than between genera, trait-based approaches may provide a first step to identify functional differences between co-occurring congeneric species in relation to mechanisms of coexistence (Tilman 1994). Therefore, the aim of the present review is to compare life history and ecological traits between some coexisting *Panulirus* species as a first step to suggesting potential trade-offs that may promote their coexistence.

There are numerous studies addressing biological and/or ecological traits of spiny lobsters but few studies comparing traits between co-occurring species. For example, the co-occurrence of multiple *Panulirus* species in tropical waters of the Indo-West Pacific has been related to a differential use of habitats of adult lobsters across environmental gradients such as depth, turbidity, coral cover, and wave action (de Bruin 1969, Berry 1971, George 1974, Pitcher 1993, Coutures and Chauvet 2003), but there is little information on the life-history traits of these particular species. Therefore, emphasis is made in this review on two pairs of co-occurring congeneric species for which relatively more information is available, one from the Eastern Central Pacific region (*Panulirus gracilis* Streets, 1871 and *P. inflatus* (Bouvier, 1895)) and the other from the Caribbean region (*P. argus* (Latreille, 1804) and *P. guttatus* (Latreille, 1804)) (Fig. 1). Throughout the text, measurements are given as mean $\pm$ 95% confidence interval unless otherwise stated.

Eastern Central Pacific: *Panulirus gracilis* and *P. inflatus*

Panulirus gracilis occurs along the continental coast and islands from Peru to Mexico, and co-occurs with *P. inflatus* along most of the Pacific coast of Mexico (Holthuis 1991) (Fig. 1). These two species are so similar that they were considered as synonyms until Holthuis and Villalobos (1961) established that they constituted separate species. During 1976–1980, the biology, ecology and fisheries of both species were concurrently studied in Zihuatanejo, Mexico, by researchers from the National




Figure 1. Geographic distribution of the two pairs of sympatric *Panulirus* species addressed in the text.

Autonomous University of Mexico. Comparative analyses of the original data from these and other studies provide insight into some differences in life history and ecological traits between these two species.

Larval and postlarval traits

Both *P. inflatus* (see Johnson and Knight 1966) and *P. gracilis* (see Báez 1983) have eleven phyllosoma stages which, according to Johnson (1971), are almost identical in almost every respect, except for the presence of a subexopodal spine on the fourth pereopod in stages five through eleven of *P. gracilis* and the absence of this spine in all stages of *P. inflatus*. However, molecular analyses have shown that this and other morphological criteria are insufficient to distinguish between phyllosomata of the two species (García-Rodríguez et al. 2008). The length of the larval phase has not been determined for either species.

Johnson (1971) described the puerulus of “*P. inflatus-gracilis*” from plankton samples collected over an area where the two species co-occur, whereas Báez (1983) described the puerulus of *P. gracilis* from samples collected in an area where only this species occurs. Both pueruli are similar in size (7.0–8.9 mm carapace length, CL) and have long, spatulated antennae, which are 2.7 times the length of the body in *P. gracilis*

(see Báez 1983) and about 2 times the length of the body in “*P. inflatus-gracilis*” (see Johnson 1971). Based on these and other minor differences, Báez (1983) suggested that the puerulus of “*P. inflatus-gracilis*” described by Johnson (1971) belonged to *P. inflatus*.

Life history strategy and benthic distribution

Panulirus gracilis occupies different types of benthic habitats, from rocky bottoms with clear water to gravel-sand bottoms near river discharges where water can be considerably turbid, whereas *P. inflatus* occurs exclusively in rocky habitats with clear waters (Briones et al. 1981, Pérez-González et al. 1992, Pérez-González 2011). Although these findings suggest that *P. inflatus* might be considered a habitat specialist and *P. gracilis* a habitat generalist, an important criterion to consider a benthic species as habitat specialist is that its postlarvae settle into the same type of habitat where the adults live, and as habitat generalist that the postlarvae are able to settle in various types of habitats (Adams et al. 2006, Adams and Ebersole 2009); however, the natural settlement habitats have not been determined for either *P. inflatus* or *P. gracilis*. However, recently settled pueruli and early benthic juveniles of both species (6–24 mm carapace length, CL) were consistently found co-occurring among the profuse biota fouling the pylons of concrete piers in Zihuatanejo, but those of *P. inflatus* persist longer in this particular habitat, which resembles the rocky habitats occupied by *P. inflatus* adults (Briones-Fourzán and Lozano-Álvarez 2013). Also, Gracia and Lozano (1980) found numerous pueruli in the stomachs of benthic catfish *Occidentarius platypogon* (Günther, 1864) (previously known as *Netuma platypogon*), which they assigned to “*P. inflatus-gracilis*” following Johnson (1971). However, it is possible that those pueruli belonged to *P. gracilis* which, unlike *P. inflatus*, dwells in the same benthic habitats as *O. platypogon* (gravel-sand and muddy bottoms).

Body size, growth rate, and fecundity

Adults of *P. inflatus* and *P. gracilis* reach a similar body size (Fig. 2A). In Zihuatanejo, *P. inflatus* has a slightly larger mean size (CL) than *P. gracilis* (Fig. 2B), but mark-recapture data showed that *P. gracilis* grows almost twice as fast as *P. inflatus* (e.g. growth rate for males, *P. gracilis*: 0.96 ± 0.08 mm CL week⁻¹, *P. inflatus*: 0.56 ± 0.10 mm CL week⁻¹, Briones-Fourzán and Lozano-Álvarez 2003) (Fig. 2C). In this same location, the size of the smallest ovigerous females ever recorded and the size at which 50% of females are ovigerous (CL₅₀) were slightly larger for *P. gracilis* (47.5 mm CL and 80.0 mm CL, respectively) than for *P. inflatus* (45.6 mm CL and 77.5 mm CL, respectively) (Weinborn 1977, Gracia 1985, Briones-Fourzán and Lozano-Álvarez 1992). Both species have an extended yearly reproductive period during which individual females can produce up to four clutches (and possible more), with embryo development taking approximately three to four weeks before hatching (Briones-Fourzán and




Figure 2. Differences in some life-history traits between *Panulirus gracilis* and *P. inflatus* from Zihuatanejo, Mexico. **A** carapace length (CL) distribution (n *P. gracilis*: 2162, n *P. inflatus*: 1873) **B** mean size **C** growth rate of males (mm CL week⁻¹, n *P. gracilis*: 148, n *P. inflatus*: 34) **D** brood size (number of eggs per clutch) versus CL relationship. Error bars denote 95% confidence intervals. (Data from **A**, **B** Briones-Fourzán and Lozano-Álvarez 1992, **C** Briones-Fourzán and Lozano-Álvarez 2003, **D** Gracia 1985, Fernández-Lomelín 1992).

Lozano-Álvarez 1992, Torres-Zepeda et al. 2008). Using original data on clutch size (number of eggs) versus CL concurrently taken for *P. inflatus* (see Gracia 1985) and *P. gracilis* (see Fernández-Lomelín 1992) from Zihuatanejo (Fig. 2D), an analysis of covariance (ANCOVA) showed that, after controlling for the significant effect of CL ($F_{1,82} = 231.71$, $p < 0.0001$), size-specific fecundity is significantly greater in *P. gracilis* than in *P. inflatus* ($F_{1,82} = 16.24$, $p < 0.001$) (Fig. 2D). Large broods are achieved partly through selection for smaller egg size. Pollock (1997) showed that, for spiny lobsters and other crustaceans, the number of eggs per gram of body weight provides an inverse index of egg size (i.e., the larger the index the smaller the egg). The use of this index shows that the eggs of *P. gracilis* (1047 ± 87 eggs g^{-1} body weight) are indeed significantly smaller than those of *P. inflatus* (911 ± 54 eggs g^{-1} body weight) (Student's t -test, $t_{85} = 2.685$, $p = 0.009$).

Use of habitat resources

Stomach content analyses showed that *P. gracilis* and *P. inflatus* consume various types of invertebrate prey but that both species exhibit a marked preference for molluscs (Lozano-Álvarez and Aramoni-Serrano 1996) (Fig. 3). In Zihuatanejo, a capture-recapture experiment was conducted during 1979–80 to estimate monthly lobster densities on a 36-ha rocky site ("site A") where the two species co-occurred (Lozano et al. 1982). At the same time, the seasonal composition of the benthic community at site A and other sites, as well as the seasonal changes in condition factor of the two lobster species were studied (Aramoni-Serrano 1982, Lozano-Álvarez and Aramoni-Serrano 1996). The total density of lobsters on site A showed a marked increase in September–October relative to the other months (Fig. 4A). For each separate species, the density showed values ≤ 15 ind. ha^{-1} between April and August, but then more than doubled in September. In October, the density of *P. inflatus* doubled again while that of *P. gracilis* decreased to previous levels. By November, the density of *P. inflatus* also decreased to previous levels (Fig. 4A).

Interestingly, site A (but not other sites) exhibited a peak in relative abundance of molluscs in the autumn (Fig. 4B), suggesting that this particular site became a food-rich habitat patch for lobsters during this season (Aramoni-Serrano 1982) and potentially explaining the substantial increase in local density of both *P. inflatus* and *P. gracilis* in September, which possibly reached the carrying capacity of the site. In October, the increase in density of *P. inflatus* and decrease in density of *P. gracilis* was followed by the recapture of several *P. gracilis* lobsters during October to December on a gravel-sand site 3 km away from site A (Lozano et al. 1982), suggesting that *P. inflatus* was able to displace *P. gracilis* from the food-rich patch. However, the high density of *P. inflatus* during the autumn possibly resulted in intraspecific competition for food resources, further exacerbated by the local decrease in abundance of molluscs during the winter (Lozano-Álvarez and Aramoni-Serrano 1996), as indicated by the significantly lower condition factor of *P. inflatus* during the winter relative




Figure 3. Diet of *Panulirus gracilis* and *P. inflatus* from Zihuatanejo, Mexico. For each food item the index of relative importance (IRI) is estimated as $IRI = (\% \text{ frequency} \times \% \text{ weight})/100$. (Data from Lozano-Álvarez and Aramoni-Serrano 1996).

to other seasons and to *P. gracilis* (Fig. 4C). In conjunction, these findings suggest that *P. inflatus* is the superior competitor in the rocky habitats to which this species is restricted.

Greater Caribbean region: *Panulirus argus* and *P. guttatus*

Panulirus argus and *P. guttatus* co-occur throughout the Greater Caribbean region (see Fig. 1). Some of the following sections are based on studies on these lobsters conducted by researchers from the National Autonomous University of Mexico in the Caribbean coast of Mexico, where life history traits and ecological aspects of these species have been concurrently studied for over 20 years.

Larval and postlarval traits

Lewis (1951) described eleven phyllosoma stages for *P. argus* from plankton samples and estimated the larval duration in about six months, but Goldstein et al. (2008), who obtained the complete larval phase in the laboratory, identified only ten distinct stages with the entire larval duration in these controlled conditions varying between 4.6 and 6.6 months. The early phyllosoma stages of *P. guttatus* have not been described and the larval duration of this species has not been determined. According to Chitty (1973),




Figure 4. Potential ecological interactions between *Panulirus gracilis* and *P. inflatus* in a rocky site ("Site A") in Zihuatanejo, Mexico. **A** lobster density (number of individuals ha⁻¹) **B** relative abundance of molluscs (percentage of molluscs in benthic samples) **C** condition factor of lobsters. Error bars denote 95% CI. (Data from **A** Lozano et al. 1982, **B** Aramoni-Serrano 1982, **C** Lozano-Álvarez and Aramoni-Serrano 1996).

first-stage phyllosomata of *P. argus* and *P. guttatus* are virtually indistinguishable based on morphology and size. However, stages VI to X of *P. guttatus* are substantially larger than the corresponding stages of *P. argus* (Baisre and Alfonso 1994).

The puerulus of *P. argus* is relatively small (6.1 mm CL on average) and has tapered antennae about 1.5 times the length of the body (Lewis et al. 1952, Goldstein et al. 2008). In contrast, the puerulus of *P. guttatus* is quite large (10 mm CL) and has long, spatulated antennae about 2.5 times the length of the body (Briones-Fourzán and McWilliam 1997). However, upon molting into the first juvenile stage and as individuals continue to grow, the antennae of *P. guttatus* become progressively shorter and thinner than those of *P. argus* (Briones-Fourzán et al. 2006).

Life history strategy and benthic distribution

It is well known that *P. argus* is an ontogenetic shifter, i.e. a species wherein the post-larvae settle into habitats distinct from those of the adults and further undergo notable ontogenetic habitat shifts toward the adult habitat (Adams and Ebersole 2009). The pueruli of *P. argus* settle in vegetated habitats of shallow reef lagoons and bays (seagrass meadows, macroalgal beds, coastal mangroves), where the early benthic juveniles remain for a few months, taking shelter among the vegetation. Eventually, juveniles outgrow the protection afforded by the vegetation and seek shelter in any crevice-type structure in or adjacent to the settlement habitats before gradually moving to the coral reef habitats where the subadults and adults dwell (Butler et al. 2006). Mating and brooding of *P. argus* occur in the reef habitat, but after embryo development – which takes from three to four weeks – is completed, the females move to deeper areas to release the phyllosoma larvae (Bertelsen 2013), which develop in oceanic waters.

Upon changing habitats, ontogenetic shifters also tend to undergo changes in behavior (Adams et al. 2006). Indeed, after their first benthic habitat shift, *P. argus* lobsters change from being asocial to being highly gregarious, with multiple individuals commonly sharing individual crevice shelters (Childress and Herrnkind 1996). In addition, *P. argus* has a highly mobile lifestyle, with movement ranges increasing with lobster size. In some locations, these movements include organized mass migrations over tens to hundreds of kilometers (Herrnkind 1969).

By contrast, *P. guttatus* is a habitat specialist, as the pueruli of this species settle directly into the coral reef habitat where the juveniles and adults also dwell (Briones-Fourzán and McWilliam 1997). Individuals of *P. guttatus* are highly sedentary, with a home range for adults of approximately 100 m in radius (Lozano-Álvarez et al. 2002). Therefore, growth, mating, brooding, and egg hatching all take place in the coral reef habitat (Briones-Fourzán and Contreras-Ortiz 1999, Negrete-Soto et al. 2002). Individuals of *P. guttatus* use reef crevices as shelters, and although small groups can share crevices, many individuals dwell solitarily, reflecting a much lower degree of gregariousness than that exhibited by *P. argus* (Briones-Fourzán 1995, Sharp et al. 1997, Briones-Fourzán and Lozano-Álvarez 2008).

Body size, growth rate, and fecundity

Adults of *P. argus* and *P. guttatus* have a very different body size (Fig. 5A). For example, in the Puerto Morelos coral reef, *P. argus* has a much larger mean size (82.3 ± 2.24 mm CL) than *P. guttatus* (59.0 ± 0.83 mm CL) (Fig. 5B) and the former species also grows much faster than the latter (weekly growth rate for males, *P. argus*: 0.91 ± 0.6 mm CL week⁻¹, *P. guttatus*: 0.26 ± 0.13 mm CL week⁻¹) (Lozano-Álvarez et al. 1991, Negrete-Soto et al. 2002) (Fig. 5C). In the same location, the largest ovigerous female of *P. guttatus* ever recorded (73.5 mm CL) was smaller than the smallest ovigerous female of *P. argus* ever recorded (75.0 mm CL). In both species, large females can produce up to four broods per year (Cruz and de León 1991, Briones-Fourzán and Contreras-Ortiz 1999), but the CL₅₀ of ovigerous females is 95.5 mm CL for *P. argus* and 59.0 mm CL for *P. guttatus* (Briones-Fourzán 1995). Due to the large interspecific difference in size, size-specific fecundity is far larger in *P. argus* than in *P. guttatus* (Fonseca-Larios and Briones-Fourzán 1998, Briones-Fourzán and Contreras-Ortiz 1999) (Fig. 5D), more so when the size of the eggs is taken into account. As Pollock (1997) had previously noted, the number of eggs per gram of body weight is significantly larger (indicating smaller eggs) in *P. argus* (689 ± 27) than in *P. guttatus* (519 ± 15) ($t_{322} = 10.925$, $p < 0.0001$).

Use of reef resources

The benthic distribution of *P. argus* and *P. guttatus* overlaps in the coral reef habitat. In Puerto Morelos, *P. guttatus* outnumbers *P. argus* by 5 to 1 across the entire reef habitat, but the relative density of each species varies with reef zone. Thus, the ratio of *P. guttatus* to *P. argus* is, on average, 2:1 in the back reef (the protected reef zone facing the mainland), but 16:1 in the fore reef (the exposed reef zone facing the open waters) (Lozano-Álvarez et al. 2007). A numerical dominance of *P. guttatus* over *P. argus* on fore reefs has also been reported in Florida (Sharp et al. 1997) and Belize (Acosta and Robertson 2003). However, there is no evidence that *P. guttatus* can displace *P. argus* via interference competition because individuals of these congeneric species do not act aggressively toward each other even when in close proximity (Lozano-Álvarez and Briones-Fourzán 2001). Rather, there is evidence that *P. guttatus* and *P. argus* make a differential use of reef resources (Lozano-Álvarez et al. 2007). For example, although lobsters of both species feed on a wide variety of organisms with a marked preference for crustaceans and molluscs (Colinas-Sánchez and Briones-Fourzán 1990) (Fig. 6), interspecific competition for food resources is unlikely, as individuals of *P. guttatus* forage on the reef itself (Wynne and Côté 2007) whereas reef-dwelling individuals of *P. argus* forage on seagrass and soft-bottom areas adjacent to the coral reefs (Cox et al. 1997, Briones-Fourzán et al. 2003).

Also, *P. argus* lobsters tend to occupy crevices ('dens') along the lower and middle portions of the reef and *P. guttatus* lobsters over the middle and upper portions of the reef (Lozano-Álvarez et al. 2007). Occupation of individual dens by multiple conspecifics is more common for *P. argus*, whereas occupation of individual dens by solitary




Figure 5. Differences in some life-history traits between *Panulirus argus* and *P. guttatus* from Puerto Morelos, Mexico. **A** carapace length (CL) distribution (n *P. argus*: 717, n *P. guttatus*: 450) **B** mean size **C** growth rate of males (mm CL week⁻¹, n *P. argus*: 148, n *P. guttatus*: 57) **D** brood size (number of eggs per clutch) versus CL relationship. Error bars denote 95% confidence intervals. (Data from **A**, **B** Lozano-Álvarez et al. 2007, Briones-Fourzán and Lozano-Álvarez 2013, **C** Negrete-Soto et al. 2002, **D** Fonseca-Larios and Briones-Fourzán 1998, Briones-Fourzán and Contreras-Ortiz 1999).




Figure 6. Diet of *Panulirus argus* and *P. guttatus* from Puerto Morelos, Mexico. For each food item the index of relative importance (IRI) is estimated as $IRI = (\% \text{ frequency} \times \% \text{ weight})/100$. (Data from Colinas-Sánchez and Briones-Fourzán 1990).

individuals is more common for *P. guttatus*. Moreover, in dens harboring either species separately or both species together, *P. argus* lobsters typically occupy the floor and the entrances of the dens, while *P. guttatus* lobster typically occupy the deep recesses of the dens, clinging to the ceiling or walls (Sharp et al. 1997, Lozano-Álvarez et al. 2007, Briones-Fourzán and Lozano-Álvarez 2008). The differential pattern of occupation of the den space by these congeners appears to minimize competition for shelter and to be related with a differential susceptibility to predators.

Susceptibility to predators

An important ecological trade-off that favors local coexistence of similar species within the same trophic level is a differential susceptibility to predators (Amarasekare 2008). This has been explored for *P. guttatus* and *P. argus*, which are potential prey to the same predators in the coral reef habitat (e.g. Randall 1967). In field studies, a negative relationship between the abundances of predators and prey would suggest that the abundance of predators controls the abundance of prey (e.g. Sih 1984, Hixon and Beets 1993, Eggleston et al. 1997). In the Puerto Morelos coral reef, Lozano-Álvarez et al. (2007) examined the relationship between the densities of predators and lobsters of each species by reef zone. A negative relationship emerged only for *P. guttatus* at the fore-reef zone, where this species was numerically dominant.

In the laboratory, individuals of *P. guttatus* and *P. argus* differed significantly in performance of several defense mechanisms expressed by spiny lobsters, indicating a differential defense strategy for each species (Briones-Fourzán et al. 2006). In particular, *P. argus* relies heavily on the formidable defense of its long, strong spiny antennae, especially when multiple individuals join in cooperative defense, whereas *P. guttatus*, which has much thinner and weaker antennae, does not express communal defensive behavior at all. *Panulirus guttatus* lobsters are also more cryptic and only emerge from their shelters to forage for short periods during the night. Individuals of *P. argus* have to traverse open areas to forage and tend to remain stationary to minimize detection when a predator approaches. If a predator attacks, a *P. argus* lobster can confront the predator, turning deftly to face it at all times, lashing and raking at the predator with its antennae. By contrast, if an individual of *P. guttatus* detects an approaching predator, it retreats backwards into the nearest available reef crevice, and if attacked, it can swim backwards in a slow but protracted bout that will effectively remove it from the visual field of the predator. In essence, *P. guttatus* exhibits a shy behavioral type and a higher susceptibility to predators, whereas *P. argus* exhibits a bold behavioral type and a lower susceptibility to predators (Briones-Fourzán et al. 2006).

Chemical ecology

The behavior of spiny lobsters is largely mediated by chemical communication (Aggio and Derby 2011). Because shelter is a limiting resource for these lobsters, individuals that are seeking shelter tend to be attracted to chemical scents released by sheltered conspecifics (“attraction cues”). On the other hand, avoiding scents from a lethally injured or freshly killed conspecific (“alarm odors”), which may signal the proximity of a predator, is a particularly effective antipredator strategy for gregarious species (Dicke and Grostal 2001). However, the degree of gregariousness varies widely among spiny lobsters (Childress 2007) and is particularly different between *P. argus* and *P. guttatus* (see Briones-Fourzán and Lozano-Álvarez 2008).

Briones-Fourzán et al. (2008) compared how individuals of *P. argus* and *P. guttatus* responded to attraction cues and alarm odors from either conspecifics or congeners. As expected, individuals of both species were significantly attracted to shelters emanating conspecific attraction cues but responded neutrally to shelters emanating congeneric attraction cues. However, individuals of *P. guttatus* responded neutrally to shelters emanating either conspecific or congeneric alarm odors, whereas individuals of *P. argus* significantly avoided shelters emanating either conspecific or congeneric alarm odors. The differential responses to alarm odors between species suggest that the cryptic defensive behavior of *P. guttatus* appears to be sufficiently adaptive to offset the need to avoid dens with conspecific (and congeneric) alarm odors, whereas learning to avoid dens with alarm odors from *P. guttatus* likely increases fitness in reef-dwelling *P. argus*, which leave their reef shelters to forage elsewhere during the night and then have to return to the reef to shelter during the day (Briones-Fourzán et al. 2008).

Discussion

The present study basically describes differences and similarities in traits between *Panulirus* species that co-occur both regionally and locally. Although just showing that species differ phenotypically or ecologically is insufficient to assign those differences to the type of trade-off necessary to promote coexistence (Siepielski and McPeck 2010), differences in morphological, physiological, ecological, and behavioral traits can help generate hypotheses regarding niche differentiation and interspecific trade-offs that influence relative fitness differences and contribute to competitive inequalities (e.g. body size, fecundity, longevity, dispersal) that may lead to coexistence, especially between congeneric species that co-occur at local scales (Tilman 1994, Amarasekare 2003, HilleRisLambers et al. 2012).

Panulirus argus and *P. guttatus* differ widely in their degree of habitat specialization and exhibit broad differences in many life history and ecological traits (e.g. larval and postlarval size and morphology, adult body size, fecundity, growth rate, movement range, behavior, susceptibility to predators) (Table 1). The large differences between *P. argus* and *P. guttatus* suggest the existence of important trade-offs leading to a stable coexistence of these two congeners. For example, although these congeners share the reef habitat, *P. guttatus* is better at exploiting shelter and food resources in this habitat, but is more susceptible to predators relative to *P. argus*. In contrast, *P. inflatus* and *P. gracilis* appear more similar in some traits (e.g. larval, postlarval, and adult size, diet) but they differ in other traits (e.g. fecundity, growth rate) and in habitat use, suggesting interspecific trade-offs that may contribute to competitive inequalities (Table 2). However, much remains to be investigated on the chemical ecology and behavioral traits of *P. gracilis* and *P. inflatus* that may influence their competitive abilities and susceptibility to predators (Table 2).

An intriguing hypothesis would be whether the substantial niche differentiation and apparent interspecific trade-offs between *P. argus* and *P. guttatus* relative to *P. gracilis* and *P. inflatus* reflect an earlier divergence of the former pair of species in the evolution of the genus. Several phylogenetic analyses (e.g. McWilliam 1995, Ptacek et al. 2001, Patek and Oakley 2003, George 2006) concur in that there are two major lineages in the radiation of *Panulirus*, with species in the first lineage representing an earlier radiation than species in the second lineage. The first lineage likely radiated from an 'argus-like ancestor' from which *P. argus* split, possibly in the Mid-Miocene (18–8 mya) (Ptacek et al. 2001, George 2006). Morphological and molecular criteria place *P. guttatus* in the first lineage as well, but the origin of this species remains uncertain as the late phyllosmata and the puerulus of *P. guttatus* exhibit the long, spatulated antennae typical of species in the second lineage, probably as a result of early divergence (McWilliam 1995, Ptacek et al. 2001, George 2006). In contrast, *P. gracilis* and *P. inflatus* clearly belong to the second major lineage and these two species constitute a single clade, with *P. inflatus* possibly splitting from *P. gracilis* as recently as the late Miocene/Pliocene (8–2 mya) (George 2006).

However, differences between the two pairs of co-occurring species due to divergence times alone would imply that speciation occurred in ecological sympatry, and at

Table 1. Summary of differences in life-history and ecological traits between *Panulirus guttatus* and *P. argus* living in sympatry in the Caribbean region.

Life history or ecological trait	<i>Panulirus guttatus</i>	<i>Panulirus argus</i>
Life-history style	Habitat specialist	Ontogenetic shifter
Larval and postlarval size	Larger	Smaller
Adult size	Smaller	Larger
Growth rate	Slower	Faster
Brood size	Smaller	Larger
Egg size	Larger	Smaller
Diet	Similar?	Similar?
Foraging habitats	Coral reef	Seagrass, rubble areas
Lifestyle	Highly sedentary	Highly mobile
Degree of gregariousness	Lower	Higher
Behavioral type	Shy	Bold
Susceptibility to predators	Higher	Lower
Response to conspecific alarm odors	Neutral	Avoidance
Response to congeneric alarm odors	Neutral	Avoidance
Competitive rank (in reef habitat)	Superior	Inferior

Table 2. Summary of differences in life history and ecological traits between *Panulirus inflatus* and *P. gracilis* living in sympatry in the Eastern Central Pacific region.

Life-history or ecological trait	<i>Panulirus inflatus</i>	<i>Panulirus gracilis</i>
Life-history strategy	Habitat specialist?	Habitat generalist?
Larval and postlarval size	Similar	Similar
Adult size	Similar	Similar
Growth rate	Slower	Faster
Brood size	Smaller	Larger
Egg size	Larger	Smaller
Diet	Similar?	Similar?
Foraging habitats	Rocky areas	Rocky + gravel-sand areas
Lifestyle	Mobile	Highly mobile
Degree of gregariousness	?	?
Susceptibility to predators	?	?
Behavioral type	?	?
Response to conspecific alarm odors	?	?
Response to congeneric alarm odors	?	?
Competitive rank	Superior?	Inferior?

least some speciation in the genus *Panulirus* appears to have been the result of vicarious events associated with major changes in oceanic currents (affecting larval dispersion) due to continental plate movements (George 2006). For example, a recent range expansion northward by *P. gracilis* into the historical range of *P. inflatus* could mean sympatry between these two species is much more recent than their time of divergence.

Moreover, if life history traits such as growth rate, size at maturity, and fecundity tend to be more similar for species that occupy more similar microhabitats, this could partially explain the greater overlap between *P. gracilis* and *P. inflatus* than between *P. argus* and *P. guttatus*. For each pair of species, these hypotheses would have to be tested either via manipulative experiments involving removal or exclusion of one species to measure its impact on the other and vice versa, or by comparing locations where both species co-occur to locations where either species is absent.

More quantitative studies are also needed to determine how much overlap in the use of food resources truly exists between co-occurring species. Spiny lobsters are omnivorous consumers, but stomach content analyses suggest that some co-occurring species prefer different types of prey (e.g. Colinas-Sánchez and Briones-Fourzán 1990). The use of stable isotope analyses (SIA) (e.g. Waddington et al. 2008) may help to better define the trophic level of co-occurring *Panulirus* species. For example, a recent study using SIA in *P. guttatus* and *P. argus* from Puerto Morelos showed that small carnivores contribute more to the diet of adult *P. guttatus* whereas small herbivores contribute more to the diet of reef-dwelling *P. argus* (Segura-García et al. unpublished data). Similarly, high through-put DNA sequencing techniques (e.g. O'Rorke et al. 2012) may help identify a potential resource partitioning between the otherwise similar phyllosoma larvae of *P. gracilis* and *P. inflatus*.

Identifying mechanisms of coexistence for congeneric species that live in sympatry is an important issue for the establishment of marine protected areas by allowing identification of species that have broad or narrow habitat requirements (McPeck 1996, Acosta and Robertson 2003). It may also provide insight into how these species could respond to climate change and other human-mediated environmental impacts such as habitat loss, degradation, and fragmentation, as well as the introduction of invasive species (McPeck 1996, HilleRisLambers et al. 2012), all of which constitute pressing issues for the shallow-water *Panulirus* species (Briones-Fourzán and Lozano-Álvarez 2013).

Acknowledgments

Support for elaborating and presenting this review at the 2013 Summer Meeting of The Crustacean Society in San José, Costa Rica, was provided by Universidad Nacional Autónoma de México and Consejo Nacional de Ciencia y Tecnología (México) through Project No. 101200. Cecilia Barradas-Ortiz produced Figure 1. Comments by two anonymous reviewers greatly helped to improve this manuscript.

References

- Acosta CA, Robertson DN (2003) Comparative spatial ecology of fished spiny lobsters *Panulirus argus* and an unfished congener *P. guttatus* in an isolated marine reserve at Glover's Reef atoll, Belize. *Coral Reefs* 22: 1–9. doi 10.1007/s00338-002-0270-5

- Adams AJ, Ebersole JP (2009) Mechanisms affecting recruitment patterns of fish and decapods in tropical coastal ecosystems. In: Nagelkerken I (Ed.) Ecological Connectivity among Tropical Coastal Ecosystems. Springer, Dordrecht, 185–228. doi: 10.1007/978-90-481-2406-0_6
- Adams AJ, Dahlgren CR, Kellison GT, Kendall MS, Layman CA, Ley JA, Nagelkerken I, Serafy JE (2006) Nursery function of tropical back-reef systems. Marine Ecology Progress Series 318: 287–301. doi: 10.3354/meps318287
- Aggio J, Derby CD (2011) Chemical communication in lobsters. In: Breithaupt T, Thiel M (Eds) Chemical Communication in Crustaceans. Springer, New York, 239–256.
- Amarasekare P (2003) Competitive coexistence in spatially structured environments: a synthesis. Ecological Letters 6:1109–1122. doi: 10.1046/j.1461-0248.2003.00530.x
- Amarasekare P (2008) Spatial dynamics of keystone predation. Journal of Animal Ecology 77: 1306–1315. doi: 10.1111/j.1365-2656.2008.01439.x
- Amarasekare P, Hoopes MF, Mouquet N, Holyoak M (2004) Mechanisms of coexistence in competitive metacommunities. The American Naturalist 164: 310–326. doi: 10.1086/422858
- Aramoni-Serrano GP (1982) Alimentación de las langostas *Panulirus inflatus* (Bouvier) y *P. gracilis* Streets en Zihuatanejo, Guerrero y su relación con el bentos. Tesis profesional, Mexico City, Mexico: Universidad Nacional Autónoma de México.
- Azovsky AI (1996) The effect of scale on congener coexistence: can molluscs and polychaetes reconcile beetles to ciliates? Oikos 77:117–126. doi: 10.2307/3545591
- Báez P (1983) Larvas phyllosoma y puerulus de la langosta verde *Panulirus gracilis* Streets, 1871 procedentes de la expedición Costa Rica 1973 (Crustacea, Decapoda, Palinuridae). Revista de Biología Marina de Valparaíso 19: 79–111.
- Baisre JA, Alfonso I (1994) Later stage larvae of *Panulirus guttatus* (Latreille, 1804) (Decapoda, Palinuridae) with notes on the identification of phyllosomata of *Panulirus* in the Caribbean sea. Crustaceana 66: 32–44. doi: 10.1163/156854094X00134
- Berry PF (1971) The spiny lobsters (Palinuridae) of the east coast of Southern Africa: distribution and ecological notes. South African Oceanographic Research Institute Investigational Reports 27: 1–23.
- Bertelsen RD (2013) Characterizing daily movements, nomadic movements, and reproductive migrations of *Panulirus argus* around the Western Sambo Ecological Reserve (Florida, USA) using acoustic telemetry. Fisheries Research 144: 91–102. doi: 10.1016/j.fishres.2012.12.008
- Bohannan BJM, Kerr B, Jessup CM, Hughes JB, Sandvik G (2002) Trade-offs and coexistence in microbial microcosms. Antonie van Leeuwenhoek 81: 107–115. doi: 10.1023/A:1020585711378
- Briones P, Lozano E, Martínez-Guerrero A, Cortés S (1981) Aspectos generales de la biología y pesca de las langostas en Zihuatanejo, Gro., México (Crustacea, Decapoda, Palinuridae). Anales del Instituto de Ciencias del Mar y Limnología Universidad Nacional Autónoma de México 8(1): 79–102.
- Briones-Fourzán P (1995) Diferencias y similitudes entre *Panulirus argus* y *P. guttatus*, dos especies de langosta comunes en el Caribe mexicano. Revista Cubana de Investigaciones Pesqueras 19(2): 14–20.

- Briones-Fourzán P, Contreras-Ortiz G (1999) Reproduction of the spiny lobster *Panulirus guttatus* (Decapoda: Palinuridae) on the Caribbean coast of Mexico. *Journal of Crustacean Biology* 19: 171–179. doi: 10.2307/1549558
- Briones-Fourzán P, Lozano-Álvarez E (1992) Aspects of the reproduction of *Panulirus inflatus* (Bouvier) and *P. gracilis* Streets (Decapoda: Palinuridae) from the Pacific coast of Mexico. *Journal of Crustacean Biology* 12: 41–50. doi: 10.2307/1548717
- Briones-Fourzán P, Lozano-Álvarez E (2003) Factors affecting growth of the spiny lobsters *Panulirus gracilis* and *Panulirus inflatus* (Decapoda: Palinuridae) in Guerrero, México. *Revista de Biología Tropical* 51: 165–174.
- Briones-Fourzán P, Lozano-Álvarez E (2008) Coexistence of congeneric spiny lobsters on coral reefs: differences in conspecific aggregation patterns and their potential antipredator benefits. *Coral Reefs* 27: 275–287. doi: 10.1007/s00338-007-0322-y
- Briones-Fourzán P, Lozano-Álvarez E (2013) Essential habitats for *Panulirus* spiny lobsters. In: Phillips BF (Ed.) *Lobsters: Biology, Management, Aquaculture and Fisheries*. Wiley-Blackwell, Oxford, 186–220.
- Briones-Fourzán P, McWilliam PS (1997) Puerulus of the spiny lobster *Panulirus guttatus* (Latreille, 1804) (Palinuridae). *Marine and Freshwater Research* 48: 699–706. doi: 10.1071/MF97130
- Briones-Fourzán P, Pérez-Ortiz M, Lozano-Álvarez E (2006) Defense mechanisms and anti-predator behavior in two sympatric species of spiny lobster, *Panulirus argus* and *P. guttatus*. *Marine Biology* 149: 227–239. doi: 10.1007/s00227-005-0191-2
- Briones-Fourzán P, Ramírez-Zaldívar E, Lozano-Álvarez E (2008) Influence of conspecific and heterospecific aggregation cues and alarm odors on shelter choice by coexisting spiny lobsters. *Biological Bulletin* 215: 182–190. doi: 10.2307/25470699
- Briones-Fourzán P, Castañeda-Fernández de Lara V, Lozano-Álvarez E, Estrada-Olivo J (2003) Feeding ecology of the three juvenile phases of the spiny lobster *Panulirus argus* in a tropical reef lagoon. *Marine Biology* 142: 855–865. doi: 10.1007/s00227-003-1013-z
- Butler MJ IV, Steneck RS, Herrnkind WF (2006) Juvenile and adult ecology. In: Phillips BF (Ed.) *Lobsters: Biology, Management, Aquaculture and Fisheries*. Blackwell, Oxford, 263–309.
- Chan TY (2010) Annotated checklist of the world's marine lobsters (Crustacea: Decapoda: Astacidea, Glypheidea, Achelata, Polychelida). *Raffles Bulletin of Zoology, Supplement* 23: 153–181.
- Chesson P (2000) Mechanisms of maintenance of species diversity. *Annual Review of Ecology and Systematics* 31: 343–366. doi: 10.1146/annurev.ecolsys.31.1.343
- Childress MJ (2007) Comparative sociobiology of spiny lobsters. In: Duffy JE, Thiel M (Eds) *Evolutionary Ecology of Social and Sexual Systems: Crustaceans as Model Organisms*. Oxford University Press, Oxford, 271–293. doi: 10.1093/acprof:oso/9780195179927.003.0013
- Childress MJ, Herrnkind WF (1996) The ontogeny of social behaviour among juvenile Caribbean spiny lobsters. *Animal Behaviour* 51: 675–687. doi: 10.1006/anbe.1996.0071
- Chitty N (1973) Aspects of the reproductive biology of the spiny lobster *Panulirus guttatus* (Latreille). MSc Thesis, University of Miami, Miami, USA.
- Cobb JS (2006) The Lobster Conferences: origin, development, and impact. *Journal of Crustacean Biology* 26: 550–554. doi: 10.1651/S-2715a.1

- Colinas-Sánchez F, Briones-Fourzán P (1990) Alimentación de las langostas *Panulirus argus* y *P. guttatus* en el Caribe mexicano. *Anales del Instituto de Ciencias del Mar y Limnología Universidad Nacional Autónoma de México* 17: 89–106.
- Coutures E, Chauvet C (2003) Study of an original lobster fishery in New Caledonia (Crustacea: Palinuridae and Scyllaridae). *Atoll Research Bulletin* 504: 1–15. doi: 10.5479/si.00775630.504.1
- Cox C, Hunt JC, Lyons WF, Davis GE (1997) Nocturnal foraging of the Caribbean spiny lobster (*Panulirus argus*) on offshore reefs of Florida, USA. *Marine and Freshwater Research* 48: 671–679. doi: 10.1071/MF97198
- Cruz R, de León ME (1991) Dinámica reproductiva de la langosta (*Panulirus argus*) en el archipiélago Cubano. *Revista de Investigaciones Marinas* 12: 234–245.
- de Bruin GHP (1969) The ecology of spiny lobsters, *Panulirus* spp., of Ceylon waters. *Bulletin of the Fisheries Research Station of Ceylon* 20: 171–189.
- Dicke M, Grostal P (2001) Chemical detection of natural enemies by arthropods: an ecological perspective. *Annual Review of Ecology and Systematics* 32: 1–23. doi: 10.1146/annurev.ecolsys.32.081501.113951
- Eggleston DB, Lipcius RN, Grover JJ (1997) Predator and shelter-size effects of coral reef fish and spiny lobster prey. *Marine Ecology Progress Series* 149: 43–59. doi: 10.3354/meps149043
- Fernández-Lomelín MP (1992) Potencial reproductivo de las langostas *Panulirus gracilis* Streets 1871 y *Panulirus inflatus* (Bouvier, 1895). Tesis profesional, Universidad Nacional Autónoma de México, Mexico City, Mexico.
- Fonseca-Larios ME, Briones-Fourzán P (1998) Fecundity of the spiny lobster *Panulirus argus* (Latreille, 1804) in the Caribbean coast of Mexico. *Bulletin of Marine Science* 63: 21–32.
- García-Rodríguez FJ, Ponce-Díaz G, Muñoz-García I, González-Armas R, Pérez-Enríquez R (2008) Mitochondrial DNA markers to identify commercial spiny lobster species (*Panulirus* spp.) from the Pacific coast of Mexico: an application on phyllosoma larvae. *Fishery Bulletin* 106: 204–212.
- George RW (1974) Coral reefs and rock lobster ecology in the Indo-west Pacific region. *Proceedings of the 2nd International Coral Reef Symposium* 1: 321–325.
- George RW (2006) Tethys Sea fragmentation and speciation of *Panulirus* spiny lobsters. *Crustaceana* 78: 1281–1309. doi: 10.1163/156854005776759780
- George RW, Main AR (1967) The evolution of spiny lobsters (Palinuridae): a study of evolution in the marine environment. *Evolution* 21: 803–820. doi: 10.2307/2406775
- Goldstein JS, Matsuda H, Takenouchi T, Butler MJ (2008) The complete development of larval Caribbean spiny lobster *Panulirus argus* (Latreille, 1804) in culture. *Journal of Crustacean Biology* 28: 306–327. doi: 10.1163/20021975-99990376
- González S, Carranza A, Scarabino F, de Mello C, Ligrone A (2011) Coexistence patterns of benthic gastropods: the genus *Buccinanops* (Nassariidae) in the inner Uruguayan continental shelf and the Río de la Plata estuary. *Pan-American Journal of Aquatic Sciences* 6: 247–259.
- Gracia A (1985) Variación estacional en la fecundidad de la langosta *Panulirus inflatus* (Bouvier, 1895) (Crustacea: Decapoda: Palinuridae). *Ciencias Marinas* 11: 7–27.

- Gracia A, Lozano E (1980) Alimentación del bagre marino *Netuma platypogon* y su importancia como indicador del reclutamiento de postlarvas de langosta (Decapoda, Palinuridae) en Guerrero, México. *Anales del Centro de Ciencias del Mar y Limnología de la Universidad Nacional Autónoma de México* 7(2): 199–206.
- Groeneveld J, Goñi R, Díaz D (2013) *Palinurus* species. In: Phillips BF (Ed.) *Lobsters: Biology, Management, Aquaculture and Fisheries*. Wiley-Blackwell, Oxford, 326–356.
- Herrnkind WF (1969) Queuing behavior of spiny lobsters. *Science* 164: 1425–1427. doi: 10.1126/science.164.3886.1425
- HilleRisLambers J, Adler PB, Harpole WS, Levine JM, Mayfield MM (2012) Rethinking community assembly through the lens of coexistence theory. *Annual Reviews of Ecology, Evolution and Systematics* 43: 227–248. doi: 10.1146/annurev-ecolsys-110411-160411
- Hixon MA, Beets JP (1993) Predation, prey refuges, and the structure of coral reef fish assemblages. *Ecological Monographs* 63: 77–101. doi: 10.2307/2937124
- Holthuis LB (1991) Marine lobsters of the world: an annotated and illustrated catalogue of species of interest to fisheries known to date. FAO Fisheries Synopsis No. 125, vol 13, FAO, Rome.
- Holthuis LB, Villalobos A (1961) *Panulirus gracilis* Streets y *Panulirus inflatus* (Bouvier), dos especies de langosta (Crustacea, Decapoda) de la costa del Pacífico de América. *Anales del Instituto de Biología Universidad Nacional Autónoma de México* 32: 251–276.
- Jeffs AG, Gardner C, Cockroft A (2013) *Jasus* and *Sagmariasus* species. In: Phillips BF (Ed.) *Lobsters: Biology, Management, Aquaculture and Fisheries*. Wiley-Blackwell, Oxford, 259–288.
- Johnson MW (1971) The palinurid and scyllarid lobster larvae from the tropical eastern Pacific and their distribution as related to the prevailing hydrography. *Bulletin of the Scripps Institution of Oceanography* 19: 1–36.
- Johnson MW, Knight M (1966) The phyllosoma larvae of the spiny lobster *Panulirus inflatus* (Bouvier). *Crustaceana* 10: 31–47. doi: 10.1163/156854066X00054
- Kneitel JM, Chase JM (2004) Trade-offs in community ecology: linking spatial scales and species coexistence. *Ecology Letters* 7: 69–80. doi: 10.1046/j.1461-0248.2003.00551.x
- Kneitel JM, Perrault D (2006) Disturbance-induced changes in community composition increase species invasion success. *Community Ecology* 7: 245–252. doi: 10.1556/Com-Ec.7.2006.2.11
- Lewis JB (1951) The phyllosoma larvae of the spiny lobster *Panulirus argus*. *Bulletin of Marine Science of the Gulf and Caribbean* 1: 89–103.
- Lewis JB, Moore HB, Babis W (1952) The post-larval stages of the spiny lobster *Panulirus argus*. *Bulletin of Marine Science of the Gulf and Caribbean* 2: 324–337.
- Lipcius RN, Eggleston DB (2000) Ecology and fishery biology of spiny lobsters. In: Phillips BF, Kittaka J (Eds) *Spiny Lobsters: Fisheries and Culture*. Blackwell, Oxford, 1–41. doi: 10.1002/9780470698808.ch
- Lozano E, Briones P, Santarelli L, Gracia A (1982) Densidad poblacional de *Panulirus gracilis* Streets y *P. inflatus* (Bouvier) (Crustacea: Palinuridae) en dos áreas cercanas a Zihuatanejo, Guerrero. *Ciencia Pesquera* 3: 61–73.

- Lozano-Álvarez E, Aramoni-Serrano G (1996) Alimentación y estado nutricional de las langostas *Panulirus inflatus* y *Panulirus gracilis* (Decapoda: Palinuridae) en Guerrero, México. *Revista de Biología Tropical* 44: 453–461.
- Lozano-Álvarez E, Briones-Fourzán P (2001) Den choice and occupation patterns of shelters by two sympatric lobster species, *Panulirus argus* and *Panulirus guttatus*, under experimental conditions. *Marine and Freshwater Research* 52: 1145–1155. doi: 10.1071/MF01093
- Lozano-Álvarez E, Briones-Fourzán P, Phillips BF (1991) Fisheries characteristics, growth, and movements of the spiny lobster *Panulirus argus* in Bahía de la Ascensión, México. *Fishery Bulletin* 89: 79–89.
- Lozano-Álvarez E, Carrasco-Zanini G, Briones-Fourzán P (2002) Homing and orientation in the spotted spiny lobster, *Panulirus guttatus* (Decapoda: Palinuridae), towards a subtidal coral reef habitat. *Crustaceana* 75: 859–873. doi: 10.1163/156854002321210712
- Lozano-Álvarez E, Briones-Fourzán P, Osorio-Arciniegas A, Negrete-Soto F, Barradas-Ortiz C (2007) Coexistence of congeneric spiny lobsters on coral reefs: differential use of shelter resources and vulnerability to predators. *Coral Reefs* 26: 361–373. doi: 10.1007/s00338-007-0207-0
- McPeck MA (1996) Trade-offs, food web structure, and the coexistence of habitat specialists and generalists. *American Naturalist* 148 (Supplement): S124–S138. doi: 10.1086/285906
- McWilliam PS (1995) Evolution in the phyllosoma and puerulus phases of the spiny lobster genus *Panulirus* White. *Journal of Crustacean Biology* 15: 542–557. doi: 10.2307/1548775
- Narwani A, Alexandrou MA, Oakley TH, Carroll IT, Cardinale BJ (2013) Experimental evidence that evolutionary relatedness does not affect the ecological mechanisms of coexistence in freshwater green algae. *Ecology Letters* 16: 1373–1381. doi: 10.1111/ele.12182
- Negrete-Soto F, Lozano-Álvarez E, Briones-Fourzán P (2002) Population dynamics of the spotted spiny lobster, *Panulirus guttatus*, in a coral reef on the Mexican Caribbean. *Journal of Shellfish Research* 21: 279–288.
- O’Rorke R, Lavery S, Chow S, Takeyama H, Tsai P, Beckley LE, Thompson PA, Waite AM, Jeffs A (2012) Determining the diet of larvae of western rock lobster (*Panulirus cygnus*) using high-throughput DNA sequencing techniques. *PLoS ONE* 7(1–10): e42757. doi: 10.1371/journal.pone.0042757
- Patek SN, Oakley TH (2003) Comparative tests of evolutionary trade-offs in a palinurid lobster acoustic system. *Evolution* 57: 2082–2100. doi: 10.1111/j.0014-3820.2003.tb00387.x
- Pérez-González R (2011) Catch composition of the spiny lobster *Panulirus gracilis* (Decapoda: Palinuridae) off the western coast of Mexico. *Latin American Journal of Aquatic Research* 39: 225–235. doi: 10.3856/vol39-issue2-fulltext-4
- Pérez-González R, Flores-Campaña LM, Nuñez-Pastén A (1992) Análisis de la distribución de tallas, captura y esfuerzo en la pesquería de las langostas *Panulirus inflatus* (Bouvier, 1895) y *P. gracilis* Streets, 1871 (Decapoda: Palinuridae) en las costas de Sinaloa, México. *Proceedings of the San Diego Society of Natural History* 15: 1–5.
- Phillips BF (2006) Perspectives. In: Phillips BF, Kittaka J (Eds) *Spiny Lobsters: Fisheries and Culture*. Blackwell, Oxford, 667–672.
- Phillips BF, Wahle RA, Ward TJ (2013) Lobsters as part of marine ecosystems—A review. In: Phillips BF (Ed.) *Lobsters: Biology, Management, Aquaculture and Fisheries*. Wiley-Blackwell, Oxford, 1–35.

- Pitcher CR (1993) Spiny lobsters. In: Wright A, Hill L (Eds) Nearshore Marine Resources of the South Pacific. Forum Fisheries Agency, Honiara, 539–607.
- Pollock DE (1997) Egg production and life-history strategies in some clawed and spiny lobster populations. *Bulletin of Marine Science* 61: 97–109.
- Ptacek MB, Sarver SK, Childress MJ, Herrnkind WF (2001) Molecular phylogeny of the spiny lobster genus *Panulirus* (Decapoda: Palinuridae). *Marine and Freshwater Research* 52: 1037–1048. doi: 10.1071/MF01070
- Randall JE (1967) Food habits of reef fishes of the West Indies. *Studies in Tropical Oceanography* 5: 665–847.
- Sfenthourakis S, Tzanatos E, Giokas S (2005) Species co-occurrence: the case of congeneric species and a causal approach to patterns of species association. *Global Ecology and Biogeography* 15: 39–49. doi: 10.1111/j.1466-822x.2005.00192.x
- Sharp WC, Hunt JH, Lyons WG (1997) Life history of the spotted spiny lobster, *Panulirus guttatus*, an obligate reef-dweller. *Marine and Freshwater Research* 48: 687–698. doi: 10.1071/MF97125
- Siepielski AM, McPeck MA (2010) On the evidence for species coexistence: a critique of the coexistence program. *Ecology* 91: 3153–3164. doi: 10.1890/10-0154.1
- Sih A (1984) The behavioral response race between predator and prey. *The American Naturalist* 123: 143–150. doi: 10.1086/284193
- Tilman D (1994) Competition and biodiversity in spatially structured habitats. *Ecology* 75: 2–16. doi: 10.2307/1939377
- Torres-Zepeda MG, Zepeda-Castillo JC, Meza-García JA, Solís-Hernández A, Villalejo-Fuerte M (2008) Reproducción de hembras de la langosta *Panulirus inflatus* (Bouvier, 1895) en el litoral del Pacífico de México. *Revista de Biología Marina y Oceanografía* 43: 129–136. doi: 10.4067/S0718-19572008000100013
- Waddington KI, Bellchambers LM, Vanderklift MA, Walker DI (2008) Western rock lobsters (*Panulirus cygnus*) in Western Australian deep coastal ecosystems (35–60 m) are more carnivorous than those in shallow coastal ecosystems. *Estuarine, Coastal and Shelf Science* 79: 114–120. doi:10.1016/j.ecss.2008.03.008
- Weinborn JA (1977) Estudio preliminar de la biología, ecología y semicultivo de los palinúridos de Zihuatanejo, Gro., México, *Panulirus gracilis* Streets y *Panulirus inflatus* (Bouvier). *Anales del Centro de Ciencias del Mar y Limnología Universidad Nacional Autónoma de México* 4: 27–78.
- Wynne SP, Côté IM (2007) Effects of habitat quality and fishing on Caribbean spotted spiny lobster populations. *Journal of Applied Ecology* 44: 488–494. doi: 10.1111/j.1365-2664.2007.01312.x